Title	Estuary Catchment Streamflow and Surface Runoff: 1975-2007	
Abstract	Catchment rainfall, runoff and evaporation were required to support the development of a new response-based estuary classification system based on dilution (from catchment runoff) and flushing characteristics. The classification system was used to stratify estuaries in NSW for the development of reference conditions and the design of a sampling program of chlorophyll a, turbidity and associated water quality parameters to assess estuary health.; Catchment rainfall, runoff and evaporation were also required to support estimation of the loads of Total Suspended Solids, Total Nitrogen and Total Phosphorus exported from estuary catchments.; Monthly time series of estimated stream flow, surface flow and base flow were modelled for 197 coastal NSW catchments. Stream flow was estimated using the 2CSalt model.	
Resource locato	r	
Data Quality Statement	Name: Data Quality Statement	
	Protocol: WWW:DOWNLOAD-1.0-httpdownload	
	Description:	
	Data quality statement for Estuary Catchment Streamflow and Surface Runoff: 1975-2007	
	Function: download	
<u>Estuary</u>	Name: Estuary Catchment Streamflow and Surface Runoff: 1975-2007	
Catchment Streamflow and	Protocol: WWW:DOWNLOAD-1.0-httpdownload	
Surface Runoff: 1975-2007	Description:	
1973 2007	Download ZIP package	
	Function: download	
Unique resource	identifier	
Code	63d4ecf7-e225-45ad-a5ec-79a0800e65cd	
Presentation form	Document digital	
Edition	1	
Dataset language	English	
Metadata standa	ard	
Name	ISO 19115	
Edition	2016	
Dataset URI	https://datasets.seed.nsw.gov.au/dataset/63d4ecf7-e225-45ad-a5ec-79a0800e65cd	
Purpose	This estuary dataset was developed under a new Monitoring, Evaluation and Reporting (MER) Program initiated by the NSW Government in 2007 to assess and better manage the health of natural resources across the State. The MER Program is in response to the NSW Natural Resources MER Strategy which has the objective of providing appropriate information for decision-making by natural resource managers.	
Status	Completed	
Spatial representation	textTable	

type				
Spatial reference	e system			
Code identifying the spatial reference system	4283			
Equivalent scale	1:None			
Topic category				

Keyword set				
keyword value	MARINE			
	MARINE-Estuaries			
	WATER-Hydrology			
	WATER-Surface			
Originating controlled vocabulary				
Title	ANZLIC Search Words			
Reference date	2008-05-16			
Geographic location				
West bounding longitude	148			
East bounding longitude	154			
North bounding latitude	-37.5			
South bounding latitude	-28			
Vertical extent information				
Minimum value	-100			
Maximum value	2228			
Coordinate reference system				
Authority code	urn:ogc:def:cs:EPSG::			
Code identifying the coordinate reference system	5711			
Temporal extent				
Begin position	1975-01-01			
End position	N/A			
Dataset reference date				
Resource maintenance				
Maintenance and update frequency	Unknown			
Contact info				
Contact position	Data Broker			
Organisation name	NSW Department of Climate Change, Energy, the Environment and Water			
Telephone number	131555			
Email address	data.broker@environment.nsw.gov.au			
Web address	https://www.nsw.gov.au/departments-and-agencies/dcceew			
Responsible party role	pointOfContact			

Lineage

Monthly time series of estimated stream flow, surface flow and base flow were modelled for 197 coastal NSW catchments. Stream flow was estimated using the 2CSalt model and this work has been described in:;; Mark Littleboy, Jon Sayers and Jocelyn Dela-Cruz (2009). Hydrological modelling of coastal catchments in New South Wales. 18th World IMACS / MODSIM Congress, Cairns, Australia 13-17 July 2009. http://mssanz.org.au/modsim09.;; Climate zones that reflect total rainfall and rainfall seasonality were defined by overlaying grids of average annual rainfall with proportion of average annual rainfall falling in winter months. For each of the 528 climate zones, daily weather data from 1956-2006 were extracted from the Queensland Department of Environment and Resource Management SILO dataset. The climate file closest to the centroid of each climate zone was obtained. The period 1956-1974 was used as a model warm-up period and results were extracted for 1975-2007.; ; 2CSalt (Stenson et al. 2005) was developed to provide water and salt inputs to regulated river models. It quantifies surface and subsurface contributions of salt and water export and predicts the impacts of land use change on water and salt export at a catchment scale. Outputs include monthly predictions of water and salt movement across several water pathways with a hillslope and alluvial groundwater store, leading to water and salt contributions to streams.

Limitations on public access

Scope dataset

DQ Completeness Commission

Effective date 2001-01-01

DQ Completeness Omission

Effective date 2001-01-01

DQ Conceptual Consistency

Effective date 1900-01-01

DQ Topological Consistency

Effective date 1900-01-01

Explanation Checked for missing attributes All attributes were checked

DQ Absolute External Positional Accuracy

Effective date 1900-01-01

DQ Non Quantitative Attribute Correctness

Effective date 1900-01-01

Responsible party

Contact position Data Broker

Organisation name NSW Department of Climate Change, Energy, the Environment and Water

Telephone number 131555

Email address <u>data.broker@environment.nsw.gov.au</u>

Web address https://www.nsw.gov.au/departments-and-agencies/dcceew

Responsible party role pointOfContact

Metadata point of contactContact positionData BrokerOrganisation nameNSW Department of Climate Change, Energy, the Environment and WaterTelephone number131555Email addressdata.broker@environment.nsw.gov.auWeb addresshttps://www.nsw.gov.au/departments-and-agencies/dcceewResponsible party rolepointOfContactMetadata date2024-02-26T13:07:07.266820

Metadata language